

MA3264 Mathematical Modelling

AY2022/23 Semester 1

Basic ODEs and Solutions

1. $M(x) - N(y)y' = 0$

(Separable) Separate the variables x and y and rewrite the equation as $\int M(x) dx = \int N(y) dy$.

2. $y' + P(x)y = Q(x)$

Multiply both sides by an **integrating factor** $\mu(x) = e^{\int P(x) dx}$:

$\mu(x)y' + \mu(x)P(x)y = \mu(x)Q(x)$

$\mu(x)y = \int \mu(x)Q(x) dx$

3. $y' + P(x)y = Q(x)y^n$

(Bernoulli) Let $z = y^{1-n}$, then $z' = (1-n)y^{-n}y'$. Hence we have:

$y^{-n}y' + P(x)y^{1-n} = Q(x)$
 $\frac{z'}{1-n} + P(x)z = Q(x)$

and use integrating factor.

4. $ay'' + by' + cy = 0$

Consider the **characteristic equation** $ax^2 + bx + c = 0$ with roots λ_1 and λ_2 :

- If $\lambda_1 \neq \lambda_2 \in \mathbb{R}$, then $y = c_1e^{\lambda_1x} + c_2e^{\lambda_2x}$.

- If $\lambda_1 = \lambda_2 \in \mathbb{R}$, then $y = (c_1 + c_2x)e^{\lambda x}$.

- If $\lambda_1 \neq \lambda_2 = \alpha \pm \beta i \in \mathbb{C}$, then $y = e^{\alpha x}(c_1 \cos \beta x + c_2 \sin \beta x)$.

5. $ay'' + by' + cy = r(x), r(x) \neq 0$

The goal is to find the **particular solution** y_p :


- If $r(x)$ is a polynomial of order n , guess $y_p(x)$ to be a n -th order polynomial.

- If $r(x)$ is in the form of $g(x)e^{kx}$, let $y_p(x) = u(x)e^{kx}$.

- If $r(x)$ is in the form of $g(x) \cos kx$ or $u(x) \sin kx$, let $z(x) = u(x)e^{ikx}$ and take $Re(z)$ or $Im(z)$.

Stability of Solutions

Harmonic Oscillation




Damped Oscillation

$mL\ddot{\theta} = -mgsin\theta - \frac{S}{L}\dot{\theta}$
 damping force
 $m\ddot{\theta} + S\dot{\theta} + \frac{mg}{L}\theta = 0$

{ Both real: **Overdamping**
 e.g. $\theta = B_1e^{-t} + B_2e^{-2t}$
 Dies rapidly to 0.
 Both complex: **Underdamping**
 e.g. $\theta = e^{-\alpha t}(B_1 \cos(\beta t) + B_2 \sin(\beta t))$
 $= Ae^{-\alpha t} \cos(\beta t - \delta)$
 "Quasi-Period"
 SHM with amplitude \downarrow with time.

Forced Oscillation



Amplitude Response Function:

$A(\omega) = \frac{F_0/m}{\sqrt{(\omega^2 - \alpha^2)^2 + (\frac{b}{m})^2 \alpha^2}}$

When $\alpha^2 = \omega^2 - \frac{b^2}{2m^2}$, \uparrow max
 $A_{resonance} = \frac{F_0/b\omega}{\sqrt{1 - (b^2/4m^2\omega^2)}}$

Conservation of Energy

Trick: $\frac{d}{dx}(\frac{1}{2}\dot{x}^2) = \dot{x} \frac{d\dot{x}}{dx} = \frac{dx}{dt} \frac{d\dot{x}}{dx} = \ddot{x}$


- Used to draw phase plane diagram (\dot{x} against x).

For SHM we have $m\ddot{x} = -kx$

$m \frac{d}{dx}(\frac{1}{2}\dot{x}^2) = -kx$
 $\frac{1}{2}m\dot{x}^2 = -\frac{1}{2}kx^2 + E$
 $E = \frac{1}{2}m\dot{x}^2 + \frac{1}{2}kx^2$
 KE PE

Population Models


Malthus' Model



Logistic Model

Assume $D = SN$ (e.g. starvation)

$\frac{dN}{dt} = BN - SN^2$ (Bernoulli)




Logistic Model with Harvesting


$\frac{dN}{dt} = (B - SN)N - E$

Case I: $E > \frac{B^2}{4S}$. Fish dies out.

Case II: $E < \frac{B^2}{4S}$.




Case II: $E = \frac{B^2}{4S}$.



Steady Growth Model

$\frac{dN}{dt} = (B_0 - \alpha \frac{dN}{dt})N - DN \approx \frac{B_0 - D}{\alpha} N$
 birth control

Model of Ants



Model of Investment

Profitability $P \equiv \frac{du/dt}{u} \rightarrow$ value of company

① Young company. All profits go back.

$\frac{du}{dt} = Pu$

② Well-established company \Rightarrow dividends

$\frac{du}{dt} = kPu$ $\frac{dw}{dt} = (1-k)Pu$
 investors

$u = Ue^{kPt}$
 $\begin{cases} k=0 & w = PUt \\ k \neq 0 & w = \frac{1}{k}(1-k)U[e^{kPt} - 1] \end{cases}$

Suppose I am in business for a fixed time T , then I pull out and start another business. Given P and T , how should I choose k ?

Define $x = kPT$, $y = \frac{w(T)}{U}$,

$y = (PT - x)(\frac{e^x - 1}{x})$

maximise borderline: $PT = 2$

If $PT > 2$, choose $k > 0$ s.t. $\frac{dy}{dx} = 0$

If $PT \leq 2$, choose $k = 0$.

System of 1st Order ODEs

Solve the general system

$$\frac{d}{dt} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \quad (\text{i.e. } \begin{cases} \frac{dx}{dt} = ax+by \\ \frac{dy}{dt} = cx+dy \end{cases})$$

$$r = \frac{1}{2} [\text{Tr}(B) \pm \sqrt{\text{Tr}(B)^2 - 4\text{Det}(B)}]$$

$$\begin{cases} r_+ & \xrightarrow{\text{eigenvector}} \vec{u}_+ \\ r_- & \xrightarrow{\text{eigenvector}} \vec{u}_- \end{cases}$$

The general solution is

$$\vec{u}(t) = C_+ e^{r_+ t} \vec{u}_+ + C_- e^{r_- t} \vec{u}_-$$




What if we have a non-homogenous equation

$$\frac{d}{dt} \begin{pmatrix} x \\ y \end{pmatrix} = B \begin{pmatrix} x \\ y \end{pmatrix} + F? \text{ An obvious particular}$$

$$\text{solution is } \begin{pmatrix} x \\ y \end{pmatrix} = -B^{-1}F.$$

Phase Plane Classification

Both r_1 and r_2 are real.



Both r_1 and r_2 are complex.



$\text{Re}[r] < 0$ $\text{Re}[r] > 0$ $\text{Re}[r] = 0$
spiral sink spiral source Centre

- Check the direction from what happen on the x -axis (sign $\left(\frac{dy}{dt}\right)$ when $x > 0$ and $y = 0$).

- Usually we consider the first quadrant.

Appendix: Common Integrals

Basic

$$\begin{aligned} \int k dx &= kx + C \\ \int x^n dx &= \frac{1}{n+1} x^{n+1} + C \\ \int \frac{1}{x} dx &= \ln|x| + C \\ \int e^x dx &= e^x + C \end{aligned}$$

Fractional

$$\begin{aligned} \int \frac{1}{ax+b} &= \frac{1}{a} \ln|ax+b| + C \\ \int \frac{1}{a^2+x^2} dx &= \frac{1}{a} \tan^{-1}\left(\frac{x}{a}\right) + C \\ \int \frac{1}{\sqrt{a^2-x^2}} dx &= \sin^{-1}\left(\frac{x}{a}\right) + C \\ \int \frac{1}{\sqrt{a^2+x^2}} dx &= \sinh^{-1}\left(\frac{x}{a}\right) + C \\ \int \frac{1}{\sqrt{x^2-a^2}} dx &= \cosh^{-1}\left(\frac{x}{a}\right) + C \\ \int \frac{1}{a^2-x^2} dx &= \frac{1}{a} \tanh^{-1}\left(\frac{x}{a}\right) + C \\ \int \frac{1}{x\sqrt{1-x^2}} dx &= -\text{sech}^{-1} x + C \\ \int \frac{1}{|x|\sqrt{1+x^2}} dx &= -\text{csch}^{-1} x + C \end{aligned}$$

Logarithmic

$$\int \ln x dx = x \ln x - x + C$$

Trigonometric

$$\begin{aligned} \int \cos x dx &= \sin x + C \\ \int \sin x dx &= -\cos x + C \\ \int \tan x dx &= \ln|\sec x| + C \\ \int \sec x dx &= \ln|\sec x + \tan x| + C \\ \int \sec^2 x dx &= \tan x + C \\ \int \sec x \tan x dx &= \sec x + C \\ \int \csc x \cot x dx &= -\csc x + C \\ \int \csc^2 x dx &= -\cot x + C \\ \int \sinh x dx &= \cosh x + C \\ \int \cosh x dx &= \sinh x + C \\ \int \text{sech}^2 x dx &= \tanh x + C \\ \int \text{csch}^2 x dx &= -\coth x + C \\ \int \text{sech} x \tanh x dx &= -\text{sech} x + C \\ \int \text{csch} x \coth x dx &= -\text{csch} x + C \end{aligned}$$

Appendix: Special Integrals

- Partial fractions
- Integration by parts: $\int u dv = uv - \int v du$
- $\int \sin^n x \cos^m x dx$:
Use trigonometric identities to convert it into $\sin^k x \cos x$ or $\cos^k x \sin x$.

Appendix: Trigonometric Identities

$$\begin{aligned} \sin, \cos: \sin^2 x + \cos^2 x &= 1 \\ \tan: \tan x &= \frac{\sin x}{\cos x} \\ \sec, \csc: \sec x &= \frac{1}{\cos x}; \csc x = \frac{1}{\sin x}; \\ \cot: \cot x &= \frac{1}{\tan x} = \frac{\cos x}{\sin x} \\ \sec^2 x - \tan^2 x &= 1; \csc^2 x - \cot^2 x = 1 \\ \sin(x+y) &= \sin x \cos y + \sin y \cos x \\ \sin 2x &= 2 \sin x \cos x \\ \sin \frac{x}{2} &= \pm \sqrt{\frac{1-\cos x}{2}} \\ \cos(x+y) &= \cos x \cos y - \sin x \sin y \\ \cos 2x &= \cos^2 x - \sin^2 x = 1 - 2 \sin^2 x = \cos^2 x - 1 \\ \cos \frac{x}{2} &= \pm \sqrt{\frac{1+\cos x}{2}} \\ \tan(x+y) &= \frac{\tan x + \tan y}{1 - \tan x \tan y} \\ \tan 2x &= \frac{2 \tan x}{1 - \tan^2 x} \\ \tan \frac{x}{2} &= \pm \sqrt{(1 - \cos x)(1 + \cos x)} \\ \sin x + \sin y &= 2 \sin \frac{x+y}{2} \cos \frac{x-y}{2} \\ \sin x \sin y &= \frac{\cos(x+y) - \cos(x-y)}{2} \\ \cos x + \cos y &= 2 \cos \frac{x+y}{2} \cos \frac{x-y}{2} \\ \cos x \cos y &= \frac{\cos(x-y) + \cos(x+y)}{2} \\ \sin x \cos y &= \frac{\sin(x+y) + \sin(x-y)}{2} \\ \sinh, \cosh: \cosh^2 x - \sinh^2 x &= 1 \\ \sinh x &= \frac{e^x - e^{-x}}{2}; \cosh x = \frac{e^x + e^{-x}}{2} \\ \tanh: \tanh x &= \frac{\sinh x}{\cosh x} \\ \text{sech} x &= \frac{1}{\cosh x} \\ \text{csch} x &= \frac{1}{\sinh x} \\ \coth: \coth x &= \frac{1}{\tanh x} \\ \tanh^2 x + \text{sech}^2 x &= 1 \\ \coth^2 x - \text{csch}^2 x &= 1 \\ \sinh(x+y) &= \sinh x \cosh y + \cosh x \sinh y \\ \sinh 2x &= 2 \sinh x \cosh x \\ \cosh(x+y) &= \cosh x \cosh y + \sinh x \sinh y \\ \cosh 2x &= \cosh^2 x + \sinh^2 x \\ \tanh 2x &= \frac{2 \tanh x}{1 + \tanh^2 x} \end{aligned}$$

[Annotated Version]

Quiz 2 Cheatsheet

MA3264 Mathematical Modelling

AY2022/23 Semester 1

Basic ODEs and Solutions

1. $M(x) - N(y)y' = 0$

(Separable) Separate the variables x and y and rewrite the equation as $\int M(x) dx = \int N(y) dy$.

2. $y' + P(x)y = Q(x)$

Multiply both sides by an Integrating factor $\mu(x) = e^{\int P(x) dx}$:

$$\mu(x)y' + \mu(x)P(x)y = \mu(x)Q(x)$$

$$\mu(x)y = \int \mu(x)Q(x) dx$$

3. $y' + P(x)y = Q(x)y^n$

(Bernoulli) Let $z = y^{1-n}$, then $z' = (1-n)y^{-n}y'$. Hence we have:

$$y^{-n}y' + P(x)y^{1-n} = Q(x)$$

$$\frac{z'}{1-n} + P(x)z = Q(x)$$

and use integrating factor

4. $ay'' + by' + cy = 0$

Consider the characteristic equation $ax^2 + bx + c = 0$ with roots λ_1 and λ_2 :

- If $\lambda_1 \neq \lambda_2 \in \mathbb{R}$, then $y = c_1e^{\lambda_1 x} + c_2e^{\lambda_2 x}$.

- If $\lambda_1 = \lambda_2 \in \mathbb{R}$, then $y = (c_1 + c_2x)e^{\lambda x}$.

- If $\lambda_1 \neq \lambda_2 = \alpha \pm \beta i \in \mathbb{C}$, then $y = e^{\alpha x}(c_1 \cos \beta x + c_2 \sin \beta x)$.

5. $ay'' + by' + cy = r(x), r(x) \neq 0$

The goal is to find the particular solution y_p :

- If $r(x)$ is a polynomial of order n , guess $y_p(x)$ to be a n -th order polynomial.

- If $r(x)$ is in the form of $g(x)e^{kx}$, let $y_p(x) = u(x)e^{kx}$.

- If $r(x)$ is in the form of $g(x) \cos kx$ or $u(x) \sin kx$, let $z(x) = u(x)e^{ikx}$ and take $\text{Re}(z)$ or $\text{Im}(z)$.

Stability of Solutions

Harmonic Oscillation

$$\frac{mL\ddot{\theta}}{ma} = -\frac{mg \sin \theta}{P}$$

$\ddot{x} = -\omega^2 x$

$\theta = 0 \rightarrow$ stable
 $\theta = \pi \rightarrow$ unstable

Damped Oscillation

$$\frac{mL\ddot{\theta}}{ma} = -\frac{mg \sin \theta}{P} - \frac{S L \dot{\theta}}{I}$$

damping force

$$m\ddot{\theta} + S\dot{\theta} + \frac{mg}{L}\theta = 0$$

Both real: Overdamping
 e.g. $\theta = B_1 e^{-\alpha t} + B_2 e^{-\beta t}$
 Dies rapidly to 0.

Both complex: Underdamping
 e.g. $\theta = e^{-\alpha t} (B_1 \cos(\beta t) + B_2 \sin(\beta t))$
 $= A e^{-\alpha t} \cos(\beta t - \delta)$
 "Quasi-period"

SHM with amplitude \downarrow with time.

Forced Oscillation

Hooke's Motor Angular Frequency $\omega = \sqrt{\frac{k}{m}}$

$$m\ddot{x} + kx = F_0 \cos \omega t$$

$x = A \cos(\omega t - \delta) + \frac{F_0/m}{\omega^2 - \omega_0^2} \cos(\omega t)$

$\frac{\omega - \omega_0}{2}$: Beat frequency
 $\omega = \omega_0$: Resonance $x = \frac{F_0/m}{2m\omega} \sin(\omega t)$

Amplitude Response Function:

$$A(\omega) = \frac{F_0/m}{\sqrt{(\omega^2 - \omega_0^2)^2 + (\frac{b}{m}\omega)^2}}$$

When $\omega^2 = \omega_0^2 = \frac{k}{m}$, \max

$$A_{\text{resonance}} = \frac{F_0/b}{\omega}$$

When $A = -\frac{F_0/m}{\omega^2 - \omega_0^2}$, $x = A(t) \sin(\frac{\omega - \omega_0}{2} t)$
 where $A(t) = \frac{2F_0/m}{\omega^2 - \omega_0^2} \sin(\frac{\omega + \omega_0}{2} t)$

Conservation of Energy

Trick $\frac{d}{dt}(\frac{1}{2}x^2) = x \frac{dx}{dt} = \frac{dx}{dt} \frac{dx}{dt} = \dot{x}^2$
 - Used to draw phase plane diagram (x against \dot{x})
 For SHM we have $m\ddot{x} = -kx$
 $\frac{d}{dt}(\frac{1}{2}kx^2) = -kx \dot{x}$
 $\frac{1}{2}m\dot{x}^2 = -\frac{1}{2}kx^2 + E$
 $E = \frac{1}{2}m\dot{x}^2 + \frac{1}{2}kx^2$

Population Models

Malthus' Model

$\frac{dN}{dt} = (B - D)N = kN$

$N(t) = N_0 e^{kt}$

Logistic Model

Assume $D = sN$ (e.g. starvation)
 $\frac{dN}{dt} = BN - sN^2$ (Bernoulli)

$N(t) = \frac{B/S}{1 + e^{-\frac{B}{S}t(\frac{B}{S} - 1)}}$



carrying capacity

Logistic Model with Harvesting

$$\frac{dN}{dt} = (B - sN)N - E$$

Case I: $E > \frac{B^2}{4s}$. Fish dies out.

Case II: $E < \frac{B^2}{4s}$.



Prepared by Tian Xian

Steady Growth Model

$$\frac{dN}{dt} = (B_0 - \alpha \frac{dN}{dt})N - DN \approx \frac{B_0 - D}{\alpha} N$$

(birth control)

Model of Ants

rate \downarrow when running out of ants

$$\frac{dx}{dt} = (\alpha + \beta x)(n - x) - \frac{sx}{r}$$

new ants attracted
 when no ants on the trail
 ants lose way

Model of Investment

Profitability $P = \frac{dU/dt}{U} \rightarrow$ value of company

① Young company. All profits go back.
 $\frac{dU}{dt} = PU$

② Well-established company \Rightarrow dividends
 $\frac{dU}{dt} = kPU$ $\frac{dU}{dt} = (k - k)PU$
 $U = U_0 e^{kPt}$ invest!

$\begin{cases} k=0 & \omega = PUt \\ k \neq 0 & \omega = \frac{1}{k}(k - k)U[e^{kPt} - 1] \end{cases}$

Suppose I am in business for a fixed time T , then I pull out and start another business. Given P and T , how should I choose k ?

Define $x = kPT$, $y = \frac{e^x - 1}{x}$

$y = (PT - x)(\frac{e^x - 1}{x})$
 maximize borderline: $PT = 2$
 If $PT > 2$, choose $k=0$ s.t. $\frac{dU}{dt} = 0$
 If $PT \leq 2$, choose $k \neq 0$.

In real life, friction cannot be ignored.

$$m\ddot{x} + b\dot{x} + kx = F_0 \cos(\omega t)$$

$$x(t) = \frac{\frac{1}{m} F_0 \cos(\omega t - \gamma)}{\sqrt{(\omega^2 - \omega_0^2)^2 + (\frac{b}{m}\omega)^2}}$$

Steady oscillation at frequency ω
 \Rightarrow Amplitude Response Function

[Annotated Version]

Qun: 2 Cheatsheet

System of 1st Order ODEs

Solve the general system

$$\frac{d}{dt} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \quad (\text{i.e. } \begin{cases} \frac{dx}{dt} = ax + by \\ \frac{dy}{dt} = cx + dy \end{cases})$$

$$r = \frac{1}{2} [\text{Tr}(B) \pm \sqrt{\text{Tr}(B)^2 - 4 \text{Det}(B)}]$$

$$\begin{cases} r_+ \text{ eigenvector } \vec{u}_+ \\ r_- \text{ eigenvector } \vec{u}_- \end{cases}$$

The general solution is

$$\vec{u}(t) = C_+ e^{r_+ t} \vec{u}_+ + C_- e^{r_- t} \vec{u}_-$$


What if we have a non-homogenous equation

$$\frac{d}{dt} \begin{pmatrix} x \\ y \end{pmatrix} = B \begin{pmatrix} x \\ y \end{pmatrix} + F$$

An obvious particular solution is $\begin{pmatrix} x \\ y \end{pmatrix} = -B^{-1}F$

Phase Plane Classification

Both r_1 and r_2 are real.



Both r_1 and r_2 are complex.



$\text{Re}(r) < 0$ Spiral sink $\text{Re}(r) > 0$ Spiral source $\text{Re}(r) = 0$ Centre

- Check the direction from what happen on the x-axis ($\text{sign}(\frac{dx}{dt})$ when $x > 0$ and $y = 0$)

- Usually we consider the first quadrant

Appendix: Common Integrals

Basic

$$\begin{aligned} \int k dx &= kx + C \\ \int x^n dx &= \frac{1}{n+1} x^{n+1} + C \\ \int \frac{1}{x} dx &= \ln|x| + C \\ \int e^x dx &= e^x + C \end{aligned}$$

Fractional

$$\begin{aligned} \int \frac{1}{ax+b} dx &= \frac{1}{a} \ln|ax+b| + C \\ \int \frac{1}{a^2 + x^2} dx &= \frac{1}{a} \tan^{-1}\left(\frac{x}{a}\right) + C \\ \int \frac{1}{\sqrt{a^2 - x^2}} dx &= \sin^{-1}\left(\frac{x}{a}\right) + C \\ \int \frac{1}{\sqrt{a^2 + x^2}} dx &= \sinh^{-1}\left(\frac{x}{a}\right) + C \\ \int \frac{1}{\sqrt{x^2 - a^2}} dx &= \cosh^{-1}\left(\frac{x}{a}\right) + C \\ \int \frac{1}{a^2 - x^2} dx &= \frac{1}{a} \tanh^{-1}\left(\frac{x}{a}\right) + C \\ \int \frac{1}{x\sqrt{1-x^2}} dx &= -\text{sech}^{-1} x + C \\ \int \frac{1}{|x|\sqrt{1+x^2}} dx &= -\text{csch}^{-1} x + C \end{aligned}$$

Logarithmic

$$\int \ln x dx = x \ln x - x + C$$

Trigonometric

$$\begin{aligned} \int \cos x dx &= \sin x + C \\ \int \sin x dx &= -\cos x + C \\ \int \tan x dx &= \ln|\sec x| + C \\ \int \sec x dx &= \ln|\sec x + \tan x| + C \\ \int \sec^2 x dx &= \tan x + C \\ \int \sec x \tan x dx &= \sec x + C \\ \int \csc x \cot x dx &= -\csc x + C \\ \int \csc^2 x dx &= -\cot x + C \\ \int \sinh x dx &= \cosh x + C \\ \int \cosh x dx &= \sinh x + C \\ \int \text{sech}^2 x dx &= \tanh x + C \\ \int \text{csch}^2 x dx &= -\text{coth} x + C \\ \int \text{sech} x \tanh x dx &= -\text{sech} x + C \\ \int \text{csch} x \coth x dx &= -\text{csch} x + C \end{aligned}$$

$$\int \tanh x dx = \ln|\cosh x| + C$$

$$\int \coth x dx = \ln|\sinh x| + C$$

Appendix: Special Integrals

- Partial fractions

- Integration by parts:

$$\int u dv = uv - \int v du$$

- $\int \sin^n x \cos^m x dx$:

Use trigonometric identities to convert it into $\sin^k x \cos x$ or $\cos^k x \sin x$.

Appendix: Trigonometric Identities

$$\sin, \cos: \sin^2 x + \cos^2 x = 1$$

$$\tan: \tan x = \frac{\sin x}{\cos x}$$

$$\sec, \csc: \sec x = \frac{1}{\cos x}; \csc x = \frac{1}{\sin x};$$

$$\cot: \cot x = \frac{\cos x}{\sin x}$$

$$\sec^2 x - \tan^2 x = 1; \csc^2 x - \cot^2 x = 1$$

$$\sin(x+y) = \sin x \cos y + \cos x \sin y$$

$$\sin 2x = 2 \sin x \cos x$$

$$\sin \frac{x}{2} = \pm \sqrt{\frac{1 - \cos x}{2}}$$

$$\cos(x+y) = \cos x \cos y - \sin x \sin y$$

$$\cos 2x = \cos^2 x - \sin^2 x = 1 - 2\sin^2 x = 2\cos^2 x - 1$$

$$\cos \frac{x}{2} = \pm \sqrt{\frac{1 + \cos x}{2}}$$

$$\tan(x+y) = \frac{\tan x + \tan y}{1 - \tan x \tan y}$$

$$\tan 2x = \frac{2 \tan x}{1 - \tan^2 x}$$

$$\tan \frac{x}{2} = \pm \sqrt{\frac{1 - \cos x}{1 + \cos x}}$$

$$\sin x + \sin y = 2 \sin \frac{x+y}{2} \cos \frac{x-y}{2}$$

$$\sin x \sin y = \frac{\cos(x-y) - \cos(x+y)}{2}$$

$$\cos x + \cos y = 2 \cos \frac{x+y}{2} \cos \frac{x-y}{2}$$

$$\cos x \cos y = \frac{\cos(x-y) + \cos(x+y)}{2}$$

$$\sin x \cos y = \frac{\sin(x+y) + \sin(x-y)}{2}$$

$$\sinh, \cosh: \cosh^2 x - \sinh^2 x = 1$$

$$\sinh x = \frac{e^x - e^{-x}}{2}; \cosh x = \frac{e^x + e^{-x}}{2}$$

$$\tanh: \tanh x = \frac{\sinh x}{\cosh x}$$

$$\text{sech} x = \frac{1}{\cosh x}$$

$$\text{csch} x = \frac{1}{\sinh x}$$

$$\coth, \text{coth} x = \frac{1}{\tanh x}$$

$$\tanh^2 x + \text{sech}^2 x = 1$$

$$\coth^2 x - \text{csch}^2 x = 1$$

$$\sinh(x+y) = \sinh x \cosh y + \cosh x \sinh y$$

$$\sinh 2x = 2 \sinh x \cosh x$$

$$\cosh(x+y) = \cosh x \cosh y + \sinh x \sinh y$$

$$\cosh 2x = \cosh^2 x + \sinh^2 x$$

$$\tanh 2x = \frac{2 \tanh x}{1 + \tanh^2 x}$$

$$\int \text{sech} x dx = \tan^{-1}(\sinh x) + C$$

$$\int \text{csch} x dx = \ln|\tanh(\frac{x}{2})| + C$$

Prepared by Tian Xiao

Romeo and Juliet

$$\begin{cases} \frac{dR}{dt} = aR & R(0) = \alpha \end{cases}$$


$$\begin{cases} \frac{dJ}{dt} = -bR & J(0) = \beta \end{cases}$$

$$\begin{cases} R(t) = \alpha \cos(\omega t) + \beta \sqrt{\frac{a}{b}} \sin(\omega t) \\ J(t) = \beta \cos(\omega t) + \alpha \sqrt{\frac{b}{a}} \sin(\omega t) \end{cases}$$

Hyperbolic Function Graphs



$$y = \cosh x = i \csc(ix)$$



$$y = \text{sech} x = \sec(ix)$$

